




# EZ-GRO Prime

CFIA REGISTERED BIOSTIMULANT

## GUARANTEED ANALYSIS

|                                      |       |
|--------------------------------------|-------|
| Indole-3-Butyric Acid (IBA).....     | 1.00% |
| 6-Furfurylaminopurine (Kinetin)..... | 0.25% |
| Salicylic Acid.....                  | 0.30% |
| Vitamin B1.....                      | 0.30% |
| Vitamin C.....                       | 0.30% |

## EFFICACY AND TRIAL DATA



**Guaranteed Minimum Analysis**  
 Indole-3-Butyric Acid (IBA).....1.00%  
 6-Furfurylaminopurine (Kinetin).....0.25%  
 Salicylic Acid.....0.30%  
 Vitamin B1.....0.30%  
 Vitamin C.....0.30%

**EZ-Gro Prime** contains IBA, Kinetin and Salicylic Acid biostimulants and plant growth regulators to help improve crop yields and reduce environmental stress.

**DANGER**

**Keep out of reach of children and animals:** Do not apply within 21 days of harvest. Do not re-enter treated areas within 4 hours of later application. Harmful if swallowed. May cause skin and respiratory irritation. May cause respiratory irritation. Wear protective gloves, clothing and respiratory protection. Do not eat and do not drink while using this product. Wash thoroughly after handling. Do not allow rinse water to contaminate streams, ponds and lakes, as water life may be endangered. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to treated areas below the high water mark. Do not connect an irrigation system (including greenhouse systems) for pesticide application to public water system. If exposed contact or feeling unwell, contact a health care provider. If swallowed, rinse mouth. If on skin, wash with plenty of water. Take off contaminated clothing and wash it before reuse. If in eyes, rinse thoroughly with water for several minutes. Remove contact lens if present and easy to do. Continue rinsing. If inhaled, move person to fresh air and hold comfortable for breathing. Store locked up in original container in a well-ventilated, cool, dry area, out of direct sunlight. Dispose of contents and packaging in accordance with local, regional and federal regulations.

**APPLICATION CHART:**

| CROP  | APPLICATION - USE ENOUGH WATER FOR COVERAGE  |
|---|--|
| Barley, Oats, Rye, Spring or Winter Wheat, Canola, Flaxseed, Mustard, Safflower, Sunflower and Corn | 3-5 leaf stage rate 175 ml/ha  |
| Legumes (Soybean, Beans, Lentils, Chickpea, Peas)   | 1st application 175 ml/ha at 3-5 leaf stage. 2nd application after cut: 175 ml/ha  |
| Alfalfa, Clover, Hay  | Apply 175 ml/ha 14-21 days after emergence. 2nd application at 5-10% bloom.  |
| Cole, Cabbage, Broccoli, Cauliflower, Brussels Sprouts  | Apply 175 ml/ha 14-21 days after emergence. 2nd application mid-season.  |
| Feeding Vegetables (Tomatoes, Peppers, Eggplant)  | Apply 175 ml/ha 14-21 days after emergence. 2nd application mid-season.  |
| Cucumbers, Watermelon, Squash   | 1st application full petal 450 ml/ha. 2nd application petal fall 450 ml/ha. 3rd application 21 days after petal fall 450 ml/ha. 4th application 50 days after petal fall 450 ml/ha.  |
| Leafy Vegetables (Lettuce, Celery, Spinach, Sugar Chard)  | 1st application full petal 450 ml/ha. 2nd application petal fall 450 ml/ha. 3rd application 21 days after petal fall 450 ml/ha. 4th application 50 days after petal fall 450 ml/ha.  |
| Apples  | 1st application pre-bloom 450 ml/ha. 2nd application petal fall 450 ml/ha. 3rd application 21 days after petal fall 450 ml/ha.   |
| Stone Fruit (Peaches, Cherries, Plums, Apricots)  | 1st application leaf out/pre-bloom 220 ml/ha. 2nd application petal fall 450 ml/ha.  |
| Grapes  | 1st application at transplant. 2nd application at 5-10% bloom. 0.9 ml EZ-Gro Prime to 1 L water (1.75 ml/ha). Apply 450 ml/ha to farmers every 3-4 weeks.  |
| Rowers & Bedding Species  | Apply 15 ml/100 ml every 2-4 weeks (only to greens & roots). Method 1 - add 1 ml/10 L water weekly through transplant, grow and bloom. Method 2 - add 1 ml/4 L water weekly throughout Bloom weeks 1-4, followed by 1/8 ml/1 L water throughout Bloom weeks 5-6 & 7. |
| Turf & Sod Production   | Apply 450 ml/ha to farmers every 3-4 weeks.  |
| Hydroponics and Medical Marijuana   | Method 1 - add 1 ml/10 L water weekly through transplant, grow and bloom. Method 2 - add 1 ml/4 L water weekly throughout Bloom weeks 1-4, followed by 1/8 ml/1 L water throughout Bloom weeks 5-6 & 7.  |

Manufactured By: Higrocorp Inc., 605 Justus Dr., Kingston, Ontario K7M 4H5 | 613-384-8882  
 Registration Number 2017100A Fertilizers Act  
 Lot No:

4 L (4.15 Kg)  
 5.6 L (5.82 Kg)

