



BIO-APP™ FE-SP TURF & FIELD WITH OSMOREGULATION

Bio-stimulant with Abiotic Plant Protection Technology.

GUARANTEED MINIMUM ANALYSIS

Fe.....6.0%
 Amino acids.....50%
 Betaine.....39%
 Monosilicic acid 0.42%
 Salicylic acid.....0.1%

FOR BEST RESULTS: foliar applications of EZ-Gro Iron (Fe) 6% 10-0-0 should be made in early morning or late evening. Some foliar burning can result if applied during hot, dry, daytime conditions. The recommended rates have been extremely successful for foliar usage, though phytotoxic symptoms may occasionally occur on sensitive crops. Consult your local agronomist for additional information on rates, timing of application and sensitivity of varieties or hybrids to phytotoxicity.

EZ-Gro Iron (Fe) 6% 10-0-0 is an Amino Acid, Chelated Micronutrient and Bio-stimulant, and is intended for use as a plant nutrient supplement for the correction of Iron deficiencies as determined by soil and/or tissue analysis. It may be used on all agricultural, horticultural and ornamental crops that require and respond to Iron nutritional supplements. EZ-Gro Iron (Fe) 6% 10-0-0 may be applied directly to the soil as a powder, through drip or injection systems, or by foliar application through conventional, low-volume, or aerial sprayers. It may be formulated into NPK fertilizers as a source of Iron, and is compatible with a wide range of fertilizer materials, pesticides, herbicides and fungicides.

MIXING INSTRUCTIONS: When making solutions of this product for soil or foliar application, DO NOT premix with water. Add this product directly to water in the solution tank, slowly and under good agitation. Continue stirring for several minutes until fully dissolved. If tank mixing with other agricultural chemicals, add this product first and allow to dissolve before adding other chemicals.

DIRECTIONS FOR USE

SOIL APPLICATION Apply to soil around plants as a powder, liquid spray, liquid injection, or through drip irrigation. When applied as a powder, this product must be watered in within 4 hours of treatment. Soil injections should be no more than three feet apart.

FOLIAR APPLICATION DO NOT apply this product when the crop is under water stress. Application after irrigation or rainfall is best. Spraying should be carried out on a calm day but not during strong sunshine. Early morning or late afternoon applications are best. If rainfall occurs within 3-4 hours after application, it may be necessary to re-spray in 3 to 4 days. If EZ-Gro Iron (Fe) 6% 10-0-0 is to be applied alone, a non-ionic wetting agent should be added to the spray tank in accordance with the manufacturer's direction. If the product is to be applied in conjunction with other chemicals which may contain their own wetting agent, then additional wetting agents may not be necessary. If using in conjunction with a hormone type selective weed killer, consult the manufacturer of the herbicide regarding the effect of wetting agents on their product. DO NOT apply to trees bearing mature fruit until you have determined that this product will not damage or mark the fruit in any way. Turf 2-4kg/ha as required, or onset of water or environmental stress, 4-5 application as required.

RATES

DEFICIENCY LEVEL	KG/ HECTARE / YEAR
Maintenance	0.5 to 1
Moderate Deficiency	1 to 1.5
Severe Deficiency	1.5 to 2

** Or as determined by tissue analysis.

ATTENTION | Keep out of reach of children and animals

This fertilizer contains iron and should be used only as recommended. It may prove harmful when misused. | Product contains soy. Adverse reactions may occur in sensitive persons. | Harmful if swallowed. | Causes skin irritation. | Causes serious eye irritation. | Wear protective clothing, gloves, a NIOSH/OSHA approved respirator, and eye protection during product mixing and handling. | Do not eat, drink or smoke while using this product. | Wash thoroughly after handling. If swallowed rinse mouth. Immediately contact a health care provider. | If in eyes, rinse thoroughly with water. Remove contact lenses if present and continue rinsing. If skin contact occurs, remove contaminated clothing and wash with soap and water. If allergic reaction occurs, seek medical attention. Wash clothing before reuse. Store locked up in original container. Dispose of contents and container in accordance with local, regional and federal regulations.



LOT NUMBER:

Manufactured By: Higrocorp Inc., 605 Justus Dr., Kingston, Ontario K7M 4H5 | 613-384-8882

Derived from Registration Numbers Fe: 2016101A | Betaine: 2017083A | Monosilicic Acid: 2017095A | Salicylic Acid: 2017070A