



## BIO-APP™ SA/SIL (FIELD)

Bio-stimulant with Abiotic Plant Protection Technology.

### GUARANTEED MINIMUM ANALYSIS

Monosilicic Acid .....14.0%  
Salicylic Acid .....6.0%

**EZ-Gro SA/SIL Field** Primes a plants SAR metabolism producing a vaccine like effects. Monosilicic acid is a plant strengthener inducing a sturdier plant along with abiotic plant protection. -Salicylic acid is a naturally occurring pgr that enhances resistance to drought, environmental stress, hardness.



### ATTENTION

#### Keep out of reach of children and animals

Causes serious skin burns and eye damage. | May be harmful if swallowed. | Do not enter treated areas within four hours of foliar application. | Wear protective gloves, clothing, eye and respiratory protection. | Do not eat, drink or smoke when using this product. | Wash thoroughly after handling. | If exposed, concerned or feeling unwell, immediately contact a health care provider. | If skin con-tact occurs, wash with soap and water. If allergic reaction occurs, seek medical attention. | If in eyes rinse cautiously with water for several minutes. Remove contact lenses if present and continue rinsing. | If swallowed rinse mouth. | Store locked up, in a dark, well ventilated area. Keep container tightly closed. | Dispose of contents and packaging in accordance with local, regional and federal legislation.

### APPLICATION CHART:

CROP	APPLICATION	TIMING/FREQUENCY
	Rate	
Nursery Stock (Containerized and field grown): deciduous and Evergreen trees, annuals, perennials, herbs, and vegetables	300ml/ha	Apply every 30 days or as needed commencing at transplant or 3-5 leaf stage.
Fruits and Berries: plums, apricots, cherry, peach, blueberries, blackberries, raspberries, strawberries	300ml/ha	3-4 applications commencing at 75% petal drop and repeating with 21-28 day intervals
Apples and Pears		3-4 applications commencing at 50% flowering and repeating with 21 day intervals
Fruiting vegetables: tomatoes, peppers, beans, peas, onions		First application prior to fruit set and repeat after 2 weeks
Lettuce, cauliflower, cabbage, Chinese cabbage		First application at 3-5 leaf stage and repeat after 2 weeks
Potatoes		First application at tuber initiation and repeat after 2 weeks
Cucurbits: pumpkins, squash, melons, cucumbers		First application at flowering and repeat after 21 days
Grapes: table and wine		Apply every 4-6 weeks commencing once plants begin to grow
Herbs: basil, chives, coriander, dill, parsley, sage		Apply every 4-6 weeks commencing once plants begin to grow
Field crops: barley, corn, oats, soybean, sugar beet, wheat		After plants reach the 3-5 leaf stage. Use with sufficient water to cover.
Grasses grown for seed, sod, pasture, forage, alfalfa		Apply when 2-4" tall. Use with sufficient water to cover.

### LOT NUMBER:

Manufactured By: Higrocorp Inc., 605 Justus Dr., Kingston, Ontario K7M 4H5 | 613-384-8882

Derived From Registration Numbers: 2017096A & 2017069A Fertilizers Act