



EZ-GRO-SA-L-10%

Guaranteed Minimum Analysis

Salicylic Acid.....	10.0%
Potassium (available K ₂ O).....	4.6%

EZ-Gro-SA-L-10% contains Salicylic Acid, a naturally occurring plant hormone, which plays a vital role in plant growth and development, photosynthesis, transpiration, ion uptake and transport. Salicylic Acid improves resistance to environmental stressors including drought, temperature, and salinity, and enhances flowering and fruit yield.

APPLICATION INSTRUCTIONS:

EZ-Gro-SA-L-10% can be used as a foliar spray, a soil drench and through fertigation systems. It can be mixed with most fungicides, insecticides and other liquid fertilizers. Do not mix concentrates together. Add water to spray tank to ¾ full followed by **EZ-Gro-SA-L-10%**, pesticides and fertilizers. | When mixing with liquid fertilizer, do not exceed the recommended rate of 50 ml/ha/application.

CAUTION

Keep out of reach of children and animals:

May be harmful if swallowed. | May cause serious eye damage. | Wear protective gloves, clothing, eye and respiratory protection during product mixing and loading. | Do not eat, drink or smoke when using this product. | Wash thoroughly after handling. | Contact a health care provider if exposed, concerned or feeling unwell. | If swallowed, rinse mouth. | If in eyes, rinse cautiously with water for several minutes. | If inhaled, move person to fresh air. | Store locked up in cool, dry, well-ventilated place out of direct sunlight. | Dispose of contents and packaging in accordance with local, regional and federal requirements.

APPLICATION CHART:

CROP	APPLICATION RATE		TIMING / FREQUENCY
	ml of product	g a.i. / ha	
Nursery Stock (Containerized and field grown): deciduous and Evergreen trees, annuals, perennials, herbs, and vegetables	40 ml / 100 L water (1.25 oz / 20 ga water)		Apply every 30 days or as needed commencing at transplant or 3-5 leaf stage.
Fruits and Berries: plums, apricots, cherry, peach, blueberries, blackberries, raspberries, strawberries			3-4 applications commencing at 75% petal drop and repeating with 21-28 day intervals
Apples and Pears			3-4 applications commencing at 50% flowering and repeating with 21 day intervals
Fruiting vegetables: tomatoes, peppers, beans, peas, onions			First application prior to fruit set and repeat after 2 weeks
Lettuce, cauliflower, cabbage, Chinese cabbage	50 ml / ha	5 g / ha	First application at 3-5 leaf stage and repeat after 2 weeks
Potatoes	(0.7 oz / acre)	(0.1 oz / acre)	First application at tuber initiation and repeat after 2 weeks
Cucurbits: pumpkins, squash, melons, cucumbers			First application at flowering and repeat after 21 days
Grapes: table and wine			Apply every 4-6 weeks commencing once plants begin to grow
Herbs: basil, chives, coriander, dill, parsley, sage			Apply every 4-6 weeks commencing once plants begin to grow
Field crops: barley, corn, oats, soybean, sugar beet, wheat			After plants reach the 3-5 leaf stage. Use with sufficient water to cover.
Grasses grown for seed, sod, pasture, forage, alfalfa			Apply when 2-4" tall. Use with sufficient water to cover.
Compost and Soil	40 ml / 100 L of soil/compost		Apply to soil or compost and mix thoroughly.
Hydroponics and Medical Marijuana	1 ml/100 L water		Apply to roots weekly through grow and bloom, beginning at transplant

- 1 L (1.4 kg)
- 3.2 L (4.6 kg)
- 6 L (8.6 kg)