



EZ-GRO SPROUT

Contains 2 Plant Growth Regulators (PGR) to Stimulate Rapid and Vigorous Root Development and Strong Plant Growth. Increases germination of seeds and promotes root growth and yields in field crops.

Guaranteed Minimum Analysis

Indole-3-Butyric Acid (IBA).....0.9%
Naphthalene Acetic Acid (NAA).....0.5%

Attention - Keep out of reach of children and animals. Store locked up in a cool, dry location. Dispose of contents and container in accordance with local, regional and national regulations.

Seed Treatment:

Apply with enough water for coverage. Compatible with most dyes

Crop	Application Rate
Soy, Cereal Grains	150 ml per metric ton of seed
Canola	75 ml per metric ton of seed

DIRECTIONS FOR USE - Apply with enough water for coverage

Field Carrots and Other Root Vegetables: Use 250 ml/1000 L water (100-150 ml/ha, 40-60 ml/acre) for foliar spray or 160 ml/ha (65 ml/acre) as in furrow. 1st application: spray the day before transplant. 2nd application: 21 days after transplant. 3rd application: 21 days after last application.

Plants, Vegetables & Herbs in Outdoor Gardens: Use 2.5 ml / 10 L water. Apply as a soil drench to thoroughly saturate roots at time of planting or when transplanting seedlings. Repeat application bi-weekly using 1.25 ml / 10 L water until plants are well established.

Plants, Vegetables & Herbs in Pots: Use 7.5 ml / 10 L water. Apply as a soil drench to thoroughly saturate roots at time of planting. Repeat application biweekly using 5 ml / 10 L water until plants are well established.

Annual & Perennial Flowers: Use 5 ml / 10 L water. Apply as a soil drench to thoroughly saturate roots at time of planting. Repeat application bi-weekly using 2.5 ml / 10 L water until plants are well established.

Propagation Benches & Beds to Reduce Transplant Shock: Use 2.5 ml / 10 L water. Thoroughly saturate roots at time of planting. Repeat application weekly using 2.5 ml / 10 L water until plants are well established.

Cuttings (Soft Wood, Semi Hard Wood, Hard Wood): Soak fresh cuttings for 30 minutes in a solution of 5 ml / 10 L water. Thoroughly saturate root area at time of planting. Repeat application weekly using 1.25 ml / 10 L water until plants are well established.

Hydroponic Greenhouse Vegetables (Cucumber, Tomato, Peppers). Use 0.5-1 ml / 10 L water every three weeks beginning at 3-5 leaf stage or transplant and continuing through grow and bloom.

Hydroponics and Medical Marijuana: Apply 0.5-1 ml/10 L water to roots, during rooting growth stage

Director - Crop Inputs Division
Directeur - Division d'intrants des cultures
Plant Health & Biosecurity Directorate
Direction de la protection et biosécurité des végétaux



NOV 15 2018

- 1 L (1.049 kg)
- 4 L (4.196 kg)