



# K

## GUARANTEED MINIMUM ANALYSIS

6-Furfurylaminopurine (Kinetin)...0.90%  
 Salicylic Acid.....0.30%  
 Vitamin B1.....0.30%  
 Vitamin C.....0.30%

**EZ-Gro K** contains biostimulants and plant growth regulators found in nature. It stimulates seedling and plant growth and may increase quality and yield. EZ-Gro K helps to reduce transplant shock and increases root weight, and the number and size of flowers.

### DANGER

#### Keep out of reach of children and animals

Do not apply within 21 days of harvest. | Do not re-enter treated areas within 12 hours of foliar application | Causes serious skin burns and eye damage | May cause respiratory irritation | Wear protective gloves, clothing, eye and respiratory protection | Do not eat, drink or smoke when using this product | Avoid breathing fumes, use only in a well-ventilated location | Wash thoroughly after handling | Do not allow rinse water to contaminate streams, ponds and lakes, as water life may be endangered | For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark | Do not connect an irrigation system (including greenhouse systems) used for pesticide application to public water system | If exposed concerned or feeling unwell, contact a health care provider | If swallowed - rinse mouth | If on skin - wash with plenty of water. Take off contaminated clothing and wash it before re-use | If in eyes - rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing | If inhaled - move person to fresh air and keep comfortable for breathing | Store locked up, tightly closed, in a dry, well-ventilated area. | Dispose of contents and packaging in accordance with local, regional, and federal requirements.

## APPLICATION CHART:

CROP	APPLICATION USE ENOUGH WATER FOR COVERAGE
Barley, Oats, Rye, Spring or Winter Wheat, Canola, Flaxseed, Mustard, Safflower, Sunflower and Corn	3-5 leaf stage rate 110 ml/ha
Legumes (Soybean, Beans, Lentils, Chickpeas, Peas)	
Alfalfa, Clover, Hay	1st application 110 ml/ha at 3-5 leaf stage. 2nd application after cut.
Kale, Cabbage, Broccoli, Cauliflower, Brussel Sprouts	
Fruiting Vegetables (Tomatoes, Peppers, Eggplant)	Apply 110 ml/ha 14-21 days after emergence. 2nd application at 5-10% bloom.
Cucumbers, Watermelon, Squash	
Leafy Vegetables (Lettuce, Celery, Spinach, Swiss Chard)	Apply 110 ml/ha 14-21 days after emergence. 2nd application mid-season.
Sugar Beets	Apply 110 ml/ha 14-21 days after emergence.
Apples	1st application full pink 290 ml/ha. 2nd application petal fall 290 ml/ha. 3rd application 21 days after petal fall 290 ml/ha. 4th application 50 days after petal fall 290 ml/ha.
Stone Fruit (Peaches, Cherries, Plums, Apricots)	1st application pre-bloom 290 ml/ha. 2nd application petal fall 290 ml/ha. 3rd application 21 days after petal fall 290 ml/ha.
Grapes	1st application leaf out/pre-bloom 140 ml/ha. 2nd application petal fall 290 ml/ha.
Flowers & Bedding Species	1st application at transplant. 2nd application at 5-10% bloom. 0.45 ml/EZ-Gro K to 1 L water (110 ml/ha).
Turf & Sod Production	Apply 8.25 ml/100 m <sup>2</sup> every 3-4 weeks foliarly to greens & tees.
Hydroponics and Cannabis	Soil/Rock Wool/Coco - Add to reservoir feed: Method 1 - add 1 ml/10 L water weekly through transplant, grow and bloom. Method 2 - add 1 ml/4 L water throughout Bloom weeks 1-4, followed by ½ ml/1 L water throughout Bloom weeks 5, 6 & 7.

## LOT NUMBER:

Manufactured By: HigoCorp Inc., 605 Justus Dr., Kingston, Ontario K7M 4H5 | 613-384-8882  
 Registration Number 2017101A Fertilizers Act

# 3.6L (3.6KG)