



BR

GUARANTEED MINIMUM ANALYSIS

24-Epibrassinolide.....0.01%

ATTENTION

Keep out of reach of children and animals

Avoid release to water sources. | Store locked up, tightly closed, in a dry, well-ventilated area. | Dispose of contents and packaging in accordance with local, regional, and federal requirements.

Brassinosteroids are naturally occurring plant hormones that promote plant growth, increase crop yield and quality, and protect against abiotic stress. **EZ-Gro BR** aids in protection against thermal, moisture, salt and heavy metal stress. **EZ-Gro BR** can optimize fertilizers by allowing the plant to use them more efficiently, thereby increasing productivity, quality and survival of crops.

APPLICATION RATES AND DIRECTIONS:

EZ-Gro BR works most effectively when used in conjunction with other fertilizers. Apply with enough water for coverage: Do not exceed the recommended rate - mg of active ingredient per hectare (mg IA/HA).

CROP	RATE (mg ai/ha)	EZ-Gro BR ml/ha	INSTRUCTIONS
General Fertigation	5	50	Apply at first signs of environmental stress
Alfalfa, clover, etc.	5	50	Apply at planting and repeat after each cutting
Beans	5	50	Apply to seeds at planting (see seed treatment) and R1 stage OR 1 application of 100 ml at 3-5 leaf stage
Canola	5	50	Apply at flowering
Cereal Grains	10	100	Apply at 3-5 leaf stage OR at boot stage
Corn, Flax	5	50	Apply at early bloom
Hay and Forage	10	100	At early active growth
Turf	10	100	Apply at spring green up and repeat every 45-60 days
Fruits, Berries	5	50	Apply at bud formation or petal fall/small fruit stage
Strawberries	10	100	Apply at spring green-up

LOT NUMBER:

continued on back

Manufactured By: HigoCorp Inc., 605 Justus Dr., Kingston, Ontario K7M 4H5 | 613-384-8882
Registration Number 2018109A Fertilizers Act

0.8L (0.85KG)

APPLICATION RATES AND DIRECTIONS CONT.:

CROP	RATE (mg ai/ha)	EZ-Gro BR ml/ha	INSTRUCTIONS
Grapes (Sizing)	30	300	Apply at small fruit size and repeat 2 weeks later
Wine Grapes	30	300	Apply at early flowering and repeat at early fruiting
Asparagus	10	100	Apply at planting, at shoot emergence, 30 days after shoot emergence
Brassicas	10	100	Apply at 50% maturity (~30 days after planting)
Cucurbits	10	100	Apply 3 weeks after planting / transplanting; optional additional treatment 3 weeks later
Leafy Annuals	10	100	Apply at planting and repeat 30 days later
Onions, Leeks, Shallots	10	100	Apply at planting and repeat 30 and 60 days later
Potatoes	10	100	Apply at tuber initiation
Sweet potatoes	10	100	Apply at planting and repeat 30-60 days later
Sugar Beets	10	100	Apply at 3 weeks post emergence
Tomatoes, peppers, eggplant	10	100	Apply shortly after planting or transplanting; repeat at 3 weeks, 6 weeks, and after first picking
Rooted Annuals	10	100	Apply at planting and repeat 3 weeks later; optional: additional treatment 6 weeks after planting
Annual Flowers	5	50	Apply at planting, at early bloom, and every 30 days during active blossom
Perennial Flowers	5	50	Apply at planting and repeat every 30 days during active growth

SOD:

Application 1	1 ml EZ-Gro BR per 1 litre of water		Spray on soil surface prior to laying sod
Application 2	15	150	Apply 2 weeks after laying sod
Application 3+	10	100	Apply every 45-60 days

SEED TREATMENT:

Dilute 1 ml EZ-Gro BR per 1 Litre of Water. Ensure adequate coverage of seeds.

HYDROPONICS AND CANNABIS:

Apply 1 ml per 10 L water to roots as injection or mixed into reservoir throughout bloom stage.

BR

