



BIO-APP™ COPPER SECURE

Bio-stimulant with Abiotic Plant Protection Technology.

GUARANTEED MINIMUM ANALYSIS

Cu.....5.0%
Salicylic Acid1.0%

Copper Salicylic complex
Bio-stimulant & Plant Protectorate
Reduces Abiotic Stress

- Primes a plants SAR metabolism producing vaccine like effects
- Supplies copper micronutrient
- Complexed with salicylic acid
- Salicylic acid reduces environmental stresses

ATTENTION

Keep out of reach of children and animals

Causes serious skin burns and eye damage. | May be harmful if swallowed. | Do not enter treated areas within four hours of foliar application. | Wear protective gloves, clothing, eye and respiratory protection. | Do not eat, drink or smoke when using this product. | Wash thoroughly after handling. | If exposed, concerned or feeling unwell, immediately contact a health care provider. | If skin con-tact occurs, wash with soap and water. If allergic reaction occurs, seek medical attention. | If in eyes rinse cautiously with water for several minutes. Remove contact lenses if present and continue rinsing. | If swallowed rinse mouth. | Store locked up, in a dark, well ventilated area. Keep container tightly closed. | Dispose of contents and packaging in accordance with local, regional and federal legislation.

APPLICATION CHART:

CROP	APPLICATION	TIMING/FREQUENCY
	Rate ml/ha	
Fruit trees (Apples, Cherries, Peaches, Pears, Blueberries etc.	150-175ml/100L water	- Beginning of flowering - 15 days after beginning petal drop - 20 days before harvest
Vegetables Greenhouse & field (Tomatoes, Peppers, Eggplants, Cucumbers, Melons, Watermelons) etc.	Fertigation 5.0L/ha Foliar: 100-125ml / 100 Liters of water	- 1 week after germination - Beginning of flowering - Small fruit stage
Leafy Vegetables, Bulb and Tuberous Vegetables (Onions, Garlic and Beets)		- 1 week after germination - Beginning of flowering/tuber formation - 30 days before harvest
Potatoes, Sweet Potato, Yams, Sugar Beet etc.		- 3-5 leaf stage - Beginning tuber initiation - 2 weeks after tuber formation
Wheat, Barley, Chickpeas, Lentils, Soy Beans, Corn		Tillering to flag leaf growth stages
Ornamentals		-Prior to flowering -At appearance of environmental stress
Cannabis (Foliar)		- 2 weeks prior to bud formation



Manufactured By: Hicrocorp Inc., 605 Justus Dr., Kingston, Ontario K7M 4H5 | 613-384-8882
Derived From Registration Number: 2017070AFertilizers Act